

ALLDES 5mg Tablets (Desloratidine)

الدرس 5 ملی گرام گولیاں
ڈیسلورٹیاڈین

DESCRIPTION:

This medicine contains an active substance Desloratidine which is an antihistamine. This is an anti allergy medicine that does not make you drowsy. It helps control your allergic reaction and its symptoms. This medicine relieves symptoms associated with allergic rhinitis (inflammation of the nasal passages caused by an allergy)

The empirical formula is C₁₇H₁₅ClN₂ and its molecular weight is 310.8 g/mol

COMPOSITION:

Each film coated tablet contains: Desloratidine.....5mg(USP Specification)

CLINICAL PHARMACOLOGY:

Pharmacokinetics

ALLDES is well absorbed from the gut and reaches highest blood plasma concentrations after about three hours. In the bloodstream, 83 to 87% of the substance are bound to plasma proteins

Mechanism of Action

ALLDES is a long-acting piperidine derivate with selective H₁ antihistaminergic and non-sedating properties. **ALLDES** diminishes the typical histaminergic effects on H₁-receptors in bronchial smooth muscle, capillaries and gastrointestinal smooth muscle, including vasodilation, bronchoconstriction, increased vascular permeability, pain, itching and spasmodic contractions of gastrointestinal smooth muscle.

INDICATION AND USAGE:

ALLDES is used to treat allergic rhinitis, nasal congestion and chronic idiopathic urticaria (hives). It is the major metabolite of loratadine and the two drugs are similar in safety and effectiveness.

An emerging indication for **ALLDES** is in the treatment of acne, as an inexpensive adjuvant to isotretinoin and possibly as maintenance therapy or monotherapy.

DOSAGE AND ADMINISTRATION:

Adults and Adolescents ≥15 years 5 mg orally once a day

Chronic idiopathic urticaria: Oral: 5 mg once daily. In one clinical trial, the titrated use of higher doses (up to 10 mg twice daily) in adults demonstrated clinical improvement .

Seasonal or perennial allergic rhinitis: Oral: 5 mg once daily

NSAID-associated urticaria, prophylaxis (off-label use): Oral: 5 mg 30 minutes before intake of a strong COX-1 inhibitor

Pediatrics

Perennial allergic rhinitis: Oral:

Infants 6 to 11 months: 1 mg once daily.

Children ≤5 years: 1.25 mg once daily.

Children 6 to 11 years: 2.5 mg once daily.

Children ≥12 years and Adolescents: 5 mg once daily.

CONTRAINDICATIONS:

Hypersensitivity to Desloratidine or to any of the excipients.

WARNINGS AND PRECAUTIONS:

Tell your doctor, pharmacist or nurse before taking this medicine:

- if you have poor kidney function

DRUG INTERACTIONS:

Alcohol (Ethyl): CNS Depressants may enhance the CNS depressant effect of Alcohol (Ethyl). *Monitor therapy*
Alizapride: May enhance the CNS depressant effect of CNS Depressants. *Monitor therapy*
Amezinium: Antihistamines may enhance the stimulatory effect of Amezinium. *Monitor therapy*
Amphetamines: May diminish the sedative effect of Antihistamines. *Monitor therapy*
Azelastine (Nasal): CNS Depressants may enhance the CNS depressant effect of Azelastine (Nasal). *Avoid combination*
Benzylpenicilloyl Polylysine: Antihistamines may diminish the diagnostic effect of Benzylpenicilloyl Polylysine. Management: Suspend systemic H1 antagonists for benzylpenicilloyl-polylysine skin testing and delay testing until systemic antihistaminic effects have dissipated. A histamine skin test may be used to assess persistent antihistaminic effects. *Consider therapy modification*

POSSIBLE SIDE EFFECTS:

Desloratadine produces potentially life-threatening effects which include Arrhythmias, Anaphylaxis, Angioedema, Extrapyramidal symptoms. which are responsible for the discontinuation of Desloratadine therapy. The symptomatic adverse reactions produced by Desloratadine are more or less tolerable and if they become severe, they can be treated symptomatically, these include Headache, Drowsiness, Fatigue, Myalgia, Palpitation, Dry mouth, Blurred vision, Urinary retention, Pharyngitis, GI disturbance, Dry mouth.

STORAGE & INSTRUCTIONS:

Keep out of reach of children & Store below 30°C.
Protect from sunlight & moisture
Improper storage may deteriorate the medicine
Do not use this medicine after the expiry date which is stated on the blister.

PRESENTATION:

ALLES 10mg film coated Tablet in pack of 10's

Manufactured by :



PHARMA LORD

PHARMA LORD | 12km, Lahore
(PVT.) LIMITED | Road, Layyah.
Pakistan
www.pharma-lord.com

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں

ہدایات: دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔

دھوپ اور نمی سے بچائیں۔ تمام ادویات بچوں کی پہنچ سے دور رکھیں

صرف رجسٹرڈ ڈاکٹر کے نسخے کے مطابق فروخت کریں۔