

Anlid

(Nimesulide)
100mg Tablets

اینلید
(نیمی سولاسیڈ)
۱۰۰ ملی گرام گولیاں

Caution:

Not to be used with impaired liver function.

Not recommended for children.

The Maximum duration of a treatment course with Nimesulide is 15 days.

Description:

Nimesulide is a nonsteroidal anti-inflammatory drug (NSAID) that selectively inhibits cyclooxygenase 2, the inducible form of the enzyme found almost exclusively in inflammatory tissues but largely devoid of physiological activity in healthy tissues, in which cyclooxygenase 1 plays important regulating roles (particularly gastric and renal physiological control).

Composition

Each film coated tablet contains: Nimesulide 100mg.

Clinical Pharmacology:

Mode of Action:

The therapeutic effects of Nimesulide are the results of its complete mode of action which targets a number of key mediators of the inflammatory process: COX-2 mediated prostaglandins, free radicals and proteolytic enzymes and histamine. Recent studies put in evidence the pharmacological activity of Nimesulide on Substance P mediator of pain in inflamed tissues. This activity may explain its effective analgesic activity.

Indications:

Nimesulide is indicated for the treatment of acute pain and primary dysmenorrhea.

Nimesulide should only be prescribed as second line treatment. The decision to prescribe Nimesulide should be based on assessment of the individual patient's overall risks.

Undesirable Effects

Gastrointestinal Disorders:

Uncommon: Gastrointestinal bleeding, Duodenal Ulcer and perforation, Gastric Ulcer and perforation.

Hepatobiliary Disorders:

Common: Hepatic enzymes increased

Possible Side Effects:

Side effects that may occur with Nimesulide are:

Uncommon: Bleeding from stomach or bowel, duodenal or stomach ulcers and burst ulcers.

Contraindications:

- Known hypersensitivity to Nimesulide or to any of the excipients of this product.
- History of hypersensitivity reactions (e.g. bronchospasm, rhinitis, urticaria) in response to acetylsalicylic acid or other non-steroidal anti-inflammatory drugs.
- History of hepatotoxic reactions to Nimesulide.
- Active gastric or duodenal ulcer, a history of recurrent ulceration or gastrointestinal bleeding, cerebrovascular bleeding or other active bleeding or bleeding disorders.
- Severe coagulation disorders.
- Severe heart failure.
- Severe renal impairment.
- Hepatic impairment.
- Children under 12 years.
- The third trimester of pregnancy and breastfeeding.

Precautions:

Symptoms/history of gastro-intestinal disease, impaired hepatic. Cardiac or renal function, pregnancy and lactation, porphyria. Caution in the elderly patients with extra cellular volume depletion from any cause. Patients on diuretics, anti-coagulants, or anti diabetics. In exceptional cases prolonged treatment proves, necessary periodic monitoring of liver function should be carried out and blood counts are recommended.

Drug interactions:

Clinically important interactions involving interference with drug metabolism have not been described with Nimesulide.

Dosage and administration:

Adults: Usual oral dose of Nimesulide in adults is 100mg tablet twice daily.

STORAGE & INSTRUCTIONS:

Keep out of reach of children & Store below 30°C.

Protect from sunlight & moisture

Improper storage may deteriorate the medicine

Presentation:

Blister pack of 20 tablets.

خوراک: ڈاکٹر ہدایت کے مطابق استعمال کریں۔
ہدایت: بچوں کی پہنچ سے دور رکھیں۔
دوا کو دھوپ، گرمی اور نمی سے محفوظ رکھیں۔ ۳۰ ڈگری سینٹی گریڈ
سے کم درجہ حرارت پر رکھیں ورنہ دوا خراب ہو جائے گی۔

Manufactured by :



PHARMA LORD | 12km, Lahore
(PVT.) LIMITED | Road, Layyah.
Pakistan
www.pharma-lord.com